The Morphine in Your Pantry – Understanding the Overdose Risk of Home-Brewed Poppy Seed Tea

Deborah Powers M.S.*,
Stephen Erickson M.D.,
Madeleine J. Swortwood Ph.D.

Department of Forensic Science Sam Houston State University Huntsville, TX, USA



Disclaimer

• No financial disclosures or conflicts of interest.

Presentation Impact


- Understand the risks involved with poppy seed tea
- Lethal amounts of morphine possible
- Reconsider the legality of unprocessed bulk poppy seeds



Opium Poppy

- Papaver somniferum
 - Grown worldwide
 - Latex harvested for alkaloids
 - Morphine
 - Codeine
 - Thebaine
 - Seeds used in baking
 - Latex transfers onto seeds during harvest

- Seeds not currently regulated
- Unwashed/unprocessed have a higher alkaloid content
- Source of morphine
 - Some codeine
- Internet culture propagates this practice




By Amazon Customer on April 5, 2017

Size: 1 LBs Verified Purchase

I use these for poppy seed tea and the trip was pretty nice.

▶ 10 comments Was this review helpful to you? Yes No Report abuse





By Ron on February 5, 2013

Verified Purchase

Ive tried a couple diff. brands of poppy seed and these are not very good. Yes they might be unwashed but if you told me they were washed i wouldn't be surprised. These did NOT do the trick, find another source.

Comment 2 people found this helpful. Was this review helpful to you? Yes No Report abuse







Case Report 1

- 24-year-old male
- Morphine intoxication with aspiration pneumonitis
- Neither illicit nor pharmaceutical sources of morphine found
- Blood sample taken at autopsy
 - Morphine: 255 ng/mL
 - Codeine: 12.2 ng/mL
 - Thebaine: detected but below the reporting limit

Case Report 2

- 21-year-old male
- Pulmonary edema and early acute pneumonia
- Neither illicit nor pharmaceutical sources of morphine found
- Blood sample taken at autopsy
 - Morphine: >800 ng/mL (femoral)
 - Codeine: 260 ng/mL (femoral)

Previous Literature

- No official LD50 for morphine or codeine in humans
 - Large overlap in therapeutic, toxic, and fatal concentrations
 - Tolerance
 - Codeine commonly prescribed in mixed formulations
- Controlled poppy seed ingestion studies
- Effects of processing on alkaloid content

Forensic Significance

- No literature on rinsing opioids by home-brewing methods
- How much morphine is possible?
 - Codeine?
- Thebaine relevant for sourcing
- Poppy seed tea abuse
 - Public safety concern



Materials

- 22 samples from internet sources
 - Bulk poppy seeds (n=19)
 - Poppy seed powder (n=1)
 - Poppy seed tea bags (n=1)
 - Liquid poppy extract (n=1)
- Analyzed for morphine, codeine, and thebaine
- Morphine-d₃ (internal standard ISTD)

Sample Inventory

Sample	Origin Claimed	Use By Date	Plant Type Claimed	Other Information
ID#	on Packaging	on Packaging	on Packaging	From Vendor
1	Canada	Nov-21	California Poppy	Poppy Seed Tea Bags
2	Unknown	Unknown	P. somniferum	Unwashed
3	Holland	Feb-18	Unknown	
4	Ohio	5/29/2017	Blue	
5	Spain	11/2018	Unknown	Unwashed
6	Unknown	Unknown	P. somniferum	
7	UK	2/16/2017	Unknown	Unwashed
8	Unknown	8/27/2018	Unknown	
9	Turkey	Jan-19	Blue	
10	England	Dec-18	Unknown	
11	Holland	May-18	Blue	Unwashed
12	Unknown	10/27/2017	Unknown	Unwashed
13	Unknown	Unknown	Black	
14	Unknown	Jun-17	Unknown	
15	Unknown	Jul-21	E. californica	Liquid Extract (400 mg/0.7 mL)
16	Tasmania	Unknown	Tasmanian Strain	Unwashed
17	Afghanistan	Unknown	Afghan Blue P. somniferum	Unwashed
18	Afghanistan	Unknown	Afghan Blue P. somniferum	Unwashed
19	USA	Oct-19	Unknown	Poppy Seed Powder
20	UK	2/11/2017	Unknown	Unwashed
21	Spain	3/28/2018	Unknown	
22	UK	8/2/2017	Unknown	Unwashed


LC-MS/MS Parameters

- Agilent 1290 Infinity II liquid chromatograph
- Agilent 6470 triple quadrupole mass spectrometer
 - Electrospray ionization (+ mode) with multiple reaction monitoring
- Agilent ZORBAX Eclipse Plus C18 column
 - 1.8 μm particle size, 2.1 x 50 mm
- Mobile Phase A: 0.1 % formic acid in water
- Mobile Phase B: 0.1 % formic acid in acetonitrile
- Flow Rate: 0.4 mL/min
- Gradient elution: hold 95:5 A:B from 0-1 min, 30 % B from 1-3 min, 95 % B from 3-3.1 min, hold until 4.4 min, decrease to 5 % B from 4.4-5 min


Sample Preparation

- 1. Tea = 85 g seeds + 150 mL water
 - 35 g powder in 100 mL, 6 g tea bags in 30 mL, and 7 mL extract [4 g] in 13 mL
- 2. Gently agitated for 10 minutes
 - room temperature neutral (RTN), room temperature acidified with 5 % lemon juice (RTA)
 - preheated to 94 °C and neutral (HN), preheated to 94 °C and acidified (HA)
- 3. Aliquots of extracts filtered
 - -Luer Lock syringes with PTFE filters
- 4. Triplicate dilutions at 1:2,000 in 95:5 (v/v) MP
 - Other dilutions when needed
- 5. Addition of ISTD at 100 ng/mL
- 6. Analyzed by validated LC-MS/MS method (SWGTOX)
 - Quantified for morphine, codeine, and thebaine from 1 − 500 ng/mL


Morphine in Seeds




Codeine in Seeds



Thebaine in Seeds



Morphine in 85 g Seeds in 150 mL Tea



Conclusion

- Alkaloid content varies
 - Between vendors
 - Between harvests
- Codeine overdose unlikely
 - About 1/10th the potency of morphine
 - Lethal amounts possible, but morphine poses greater risk
- Moderate volumes

 Lethal morphine
 - Dependent on individual tolerances
 - However, no guarantee of safety
- Heating & acid modifier could increase alkaloid content in tea

JOURNAL OF FORENSIC SCIENCES



J Forensic Sci, 2017 doi: 10.1111/1556-4029.13664 Available online at: onlinelibrary.wiley.com

PAPER

TOXICOLOGY

Deborah Powers, ¹ M.S.; Stephen Erickson, ² M.D.; and Madeleine J. Swortwood, ¹ Ph.D.

Quantification of Morphine, Codeine, and Thebaine in Home-Brewed Poppy Seed Tea by LC-MS/MS

Acknowledgements

- Sam Houston State University Department of Forensic Science
- Dr. Gayle Suzuki, Virginia Office of the Chief Medical Examiner
- Dr. Stephen Erickson, Office of the Chief Medical Examiner, Arkansas State Crime Lab
- The Hacala Family



Questions

Deborah Powers

deborah.powers.42790@gmail.com

Dr. Madeleine J. Swortwood

mjs079@shsu.edu

