

Synthetic Cathinone Stability in Blood Using LC/Q-TOF-MS

Lindsay Glicksberg, BS; Sarah Kerrigan, PhD

*Department of Forensic Science
Sam Houston State University
Huntsville, TX, USA*

69th Annual AAFS, New Orleans, 2017



**Sam Houston
State University**



FSF Emerging Forensic Scientist Award
Paper Presentation



Disclosure

- There is no real or apparent conflicts of interest related to the content of this presentation
- Products used:
 - Agilent Technologies 6530 Accurate-Mass Q-TOF LC/MS
- The authors declare no competing interest
- Project funded by NIH Award # 2013-R2-CX-K006

Synthetic Cathinone Background

- Derived from cathinone
 - *Catha edulis* shrub
- Synthesized for effects similar to methamphetamine & MDMA
- Available on the internet and in head shops with labels such as:
 - “not for human consumption”
 - “bath salts”
 - “plant food”



Pharmacology and Toxicology

Desired Effects:

- stimulant and euphoric symptoms
 - Increased energy, mood enhancement, empathy, sociability, concentration, euphoria

Adverse Effects:

- neurological, cardiovascular, and psychopathological symptoms
 - Hallucinations, delusions, confusion, violence, homicidal tendencies, death


Antemortem Cases: Motor Vehicle Accidents & Driving While Impaired

Postmortem Cases: Overdose, Suicide, Homicide



General Structure of Cathinone

- Phenethylamines
- R groups represent positions that can be substituted to create various synthetic cathinones



Importance of Stability

- Understanding the stability of a drug in biological matrices is essential
- Condition and length of storage can affect drug concentration
- Specimens stored for days, weeks, or months prior to analysis
- Subjected to various conditions during collection and shipping process

Cathinone Instability

Plant Material

- Unstable in oxygen rich conditions (Szendrei, 1980)
- Unstable in alkaline conditions (Szendrei, 1980; Berrang, 1982)
- Dimer formation (Berrang, 1982; Chappell, 2010)
- Best to air dry and refrigerate

Thermal Degradation

- Thermal degradation in GC-MS
 - Methcathinone (DeRuiter, 1994)
 - α -PVP (Tsujikawa, 2013)
 - 19 synthetic cathinones (Kerrigan, 2015)
- **Aqueous Solution (Tsujikawa, 2012)**
 - Stable at acidic pH
 - Decomposition rate dependent upon chemical structure
 - 5 synthetic cathinones

Biological Material

■ Sorensen (2011):

- Methcathinone, Ethcathinone, Mephedrone, Flephedrone, Methedrone, Methylone, Butylone
- Blood (pH 7.4 and 5.9)
- 5°C and 20°C
- 7 days
- More stable in pH 5.9 and 5°C over 7 day period

■ Johnson and Botch-Jones (2013):

- MDPV/Mephedrone
- Blood, Plasma, Urine
- -20°C, 4°C, 22°C
- 14 days
- -20°C: stable in 3 matrices
- Mephedrone unstable at 4°C and 22°C

■ Soh and Elliott (2014):

- 4-MEC
- Blood and Plasma
- 20°C and 5°C
- 7 days
- Unstable at both temperature

■ Busardo (2016):

- Mephedrone
- Antemortem & Postmortem Blood
- -20°C, 4°C, 20°C
- 6 months
- Unstable at 4°C and 20°C by 3 months
- Stable at -20°C

Stability Studies

No systematic and fully comprehensive study addressing synthetic cathinone stability in biological evidence

22 synthetic cathinones

2 biological matrix


4 temperatures

>6 months

Comprehensive study assessing stability as it relates to

1. Concentration
2. Temperature
3. Storage Time
4. Chemical Structure

Research Design



LC/Q-TOF-MS Conditions

Agilent Technologies 6530 Accurate-Mass Q-TOF LC/MS

LC Separation

- Poroshell 120 EC-C18 Column (2.1x100mm, 2.7 μ m particle size)
- Mobile Phase A: 0.1% FA in diH₂O
- Mobile Phase B: 0.1% FA in ACN
- Flow Rate: 0.40 mL/min
- LC Gradient:
 - 96% A to 5 min, 90% A until 11 min, 60% A for 1 min, 0% A to equilibrate the column

Q/TOF Parameters

- Gas Temperature: 200°C
- Gas Flow Rate: 13 L/min
- Sheath Gas Temperature: 250°C
- Sheath Gas Flow Rate: 12 L/min
- Nebulizer Pressure 20 psig


Mass Spectrometry

- Capillary Voltage: 4000 V
- Fragmentor Voltage: 150 V
- Nozzle Voltage: 0 V
- Collision Energy: 30 eV, 20 eV
- MS Scan Rate: 8 spectra/sec
- MS/MS Scan Rate: 3 spectra/sec
- MS Scan Range: 40-1000 m/z
- ESI Mode: Positive

Acquisition

- Minimum of two ion transitions per drug
- Run Time: 13 minutes

LC/Q-TOF EIC



Validation Summary

SWGTOX Standard Practices for Method Validation

LOD: 1 – 5 ng/mL

LOQ: 1 – 5 ng/mL

Inter-assay Precision: 3 – 12%

Intra-assay Precision: 0 – 14%

Bias: -7 – 11%

Accuracy: 93 – 100%

Matrix Effects: -15 – 3%

Dilution Integrity: 2- and 4- fold

Interferences: No interferences (>50 interferents)

Glücksberg, L., Bryand, K., Kerrigan, S., 2016. Identification and quantification of synthetic cathinones in blood and urine using liquid chromatography-quadrupole/time of flight (LC-Q/TOF) mass spectrometry. Journal of Chromatography B 1035, 91-103.

Stability Study Analysis

Extraction

- Blood samples in duplicate (n=2)
 - 1000 ng/mL samples 1:4 dilution
- Calibrators extracted with every run
 - 10, 25, 100, 250, 350, and 500 ng/mL
- Negative and Positive (100 ng/mL) Controls


Analysis

- Concentration Mean (n=2)
- Error bars emitted for clarity
- Significant >20% loss


Month	Samplings/Week
1	4
2-3	2-3*
4-6	1
6-12	1/month

Secondary Amines, No Ring Substituents


Methcathinone




Ethcathinone



Buphedrone




Pentedrone




Secondary Amines, Ring Substituted


Mephedrone*




4-MEC




4-EMC




Methedrone




Flephedrone



3-FMC




3,4-DMMC




Secondary Amines, Methylenedioxy-Type


Methylone*




Ethylone*




Butylone*



Pentylone*




Eutylone*




Tertiary Amines, Pyrrolidine-Type


Alpha-PVP*




MPBP




Pyrovalerone




Naphyrone*




MDPV*



MDPBP




Chemical Structure Dependence:




Temperature Dependence (100 ng/mL)

21


32°C




20°C




4°C



-20°C



Concentration Dependence




Stability Range (in days)

Cathinone Structural Group	32°C	20°C	4°C	-20°C
Unsubstituted	<1	2 – 5	19 – 34	76 – 146
Ring Substituted	<1	<1 – 7	3 – 88	14 - >184
Methylenedioxy	1 – 3	9 – 24	88 - >184	>184
Pyrrolidine	2 – 8	8 – 34	115 - >184	88 - >184

Stability Range

24



Detection Window (in days)

Cathinone Structural Group	32°C	20°C	4°C	-20°C
Unsubstituted	3 – 4	19 - 24	≥184	>184
Ring Substituted	2 – 7	7 – 55	88 - >184	>184
Methylenedioxy	14 – 27	76 – 184	>184	>184
Pyrrolidine	11 – 115	55 - >184	>184	>184

Conclusions

- **Temperature Dependence**
 $-20^{\circ}\text{C} > 4^{\circ}\text{C} > 20^{\circ}\text{C} > 32^{\circ}\text{C}$
- **No Concentration Dependence**
- **Significant Structural Dependence**
 $\text{MD/PYR} > \text{PYR} > \text{MD} > \text{Ring Substituted} > \text{Unsubstituted} > \text{3-FMC}$

Conclusions Cont.

- **Significant loss on the order of hours**


3-FMC: 32°C—undetectable after 24 hours

Substituted and Unsubstituted

32°C: >20% loss at 48 hours

20°C: >20% loss at 48 hours

- **Instability at common storage conditions (4°C)**



- **Significant Structural Influence**

Acknowledgements

This project was supported by Award No. 2013-R2-CX-K006 awarded by the National Institute of Justice, Office of Justice Programs, U.S. Department of Justice. The opinions, findings, and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect those of the Department of Justice.



FSF Emerging Forensic Scientist Award
Paper Presentation

Questions?

Lindsay Glicksberg
lcg012@shsu.edu

